

## AMERICAN SOCIETY FOR MICROBIOLOGY

## Draft Genome Sequence of *Enterobacter cloacae* 3F11 (Phylum *Proteobacteria*)

Christopher R. Dumigan,<sup>a</sup> Gregory E. Perry,<sup>a</sup> K. Peter Pauls,<sup>a</sup> Manish N. Raizada<sup>a</sup>

<sup>a</sup>Department of Plant Agriculture, University of Guelph, Guelph, Ontario, Canada

**ABSTRACT** Presented here is the draft genome sequence of *Enterobacter cloacae* 3F11. This seed endophyte solubilizes rock phosphate and was isolated from *Zea nicaraguensis*. The genome contains 4,579,108 bp in 264 contigs.

The insolubility of rock phosphate presents a major challenge for plants to obtain this vital macronutrient (1). Certain forms of rock phosphate can be made bioavailable by acidification, and some plant species have evolved organic acid excretion as an adaptation to soils rich in rock phosphate (2). Plants can also adapt to their environment by hosting nonpathogenic microbes known as endophytes that provide a selective advantage to their host (3).

A recent study identified an endophytic *Enterobacter* sp. in *Zea nicaraguensis*, a wild maize growing in a swamp at the base of the San Cristobal Volcano in Nicaragua (4). The primary form of phosphate in volcanic rock is highly insoluble. This endophytic *Enterobacter* species was found to localize to the root hair cells of modern corn and annual ryegrass (a model grass relative of corn), increase the root growth of annual ryegrass on insoluble rock phosphate, and secrete acids to solubilize rock phosphate (4).

Here, we present the whole-genome sequence of *Enterobacter* sp. strain 3F11, which was isolated from surface-sterilized seeds of *Zea nicaraguensis* in 2011 (5). Genomic DNA was isolated from a 30°C (200 rpm) overnight lysogeny broth (LB) culture grown from a single colony using a Norgen bacterial genomic DNA isolation kit (catalog number 17900; Thorold). A paired-end library was created using a Nextera XT DNA library prep kit (Illumina, San Diego, CA), and sequencing was performed on an Illumina MiSeq platform. The analysis generated 1,298,802 read pairs averaging 151 bp in size. The total genome coverage was 43-fold, and 1,297,520 reads remained after trimming low-quality reads. *De novo* assembly was performed using CLC Genomics Workbench 10.0.1 (Qiagen) and yielded a genome size of 4,579,108 bp in 264 contigs (minimum, 1,059 bp; minimum, 124,272 bp;  $N_{50}$ , 29,224 bp), with a GC content of 56.4%.

Genome annotation, carried out by the RAST server (6), identified 4,201 proteinencoding genes, 58 tRNA operons, and 3 rRNA operons. Taxonomy was determined using K-mer finder (7, 8), and the closest match was found to be *Enterobacter cloacae*, with 98.6% nucleotide similarity to Hoffman complex IV (9); thus, we propose the strain name *E. cloacae* IV 3F11.

**Data availability.** The results of the whole-genome shotgun project were deposited at DDBJ/EMBL/GenBank under the accession number QKVR00000000. The version described in this paper is version QKVR01000000. Raw Illumina reads are available at SRA accession number SRR7310671.

## ACKNOWLEDGMENTS

We thank David Johnston-Monje for the original isolation of this strain while he was a Ph.D. student at the University of Guelph.

This research was supported by a LAAIR grant to M.N.R. jointly funded by The

Received 26 June 2018 Accepted 12 July 2018 Published 16 August 2018

Citation Dumigan CR, Perry GE, Pauls KP, Raizada MN. 2018. Draft genome sequence of *Enterobacter cloacae* 3F11 (phylum *Proteobacteria*). Microbiol Resour Announc 7:e00846-18. https://doi.org/10.1128/MRA .00846-18.

**Editor** Catherine Putonti, Loyola University Chicago

**Copyright** © 2018 Dumigan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Manish N. Raizada, raizada@uoguelph.ca.

Ontario Ministry of Agriculture, Food and Rural Affairs and The Canadian First Research Excellent Fund, a Canadian Foundation for Innovation grant to K.P.P., and a Queen Elizabeth II Graduate Scholarship to C.R.D. from the Government of Canada.

## REFERENCES

- 1. Hinsinger P. 2001. Bioavailability of soil inorganic P in the rhizosphere as affected by root-induced chemical changes: a review. Plant Soil 237: 173–195. https://doi.org/10.1023/A:1013351617532.
- Dakora FD, Phillips DA. 2002. Root exudates as mediators of mineral acquisition in low-nutrient environments. Plant Soil 245:35–47. https:// doi.org/10.1023/A:1020809400075.
- Rosenblueth M, Martínez-Romero E. 2006. Bacterial endophytes and their interactions with hosts. Mol Plant Microbe Interact 19:827–837. https:// doi.org/10.1094/MPMI-19-0827.
- Shehata HR, Dumigan C, Watts S, Raizada MN. 2017. An endophytic microbe from an unusual volcanic swamp corn seeks and inhabits root hair cells to extract rock phosphate. Sci Rep 7:13479. https://doi.org/10 .1038/s41598-017-14080-x.
- Johnston-Monje D, Raizada MN. 2011. Conservation and diversity of seed associated endophytes in *Zea* across boundaries of evolution, ethnography and ecology. PLoS One 6:e20396. https://doi.org/10.1371/journal .pone.0020396.
- 6. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K,

Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. BMC Genomics 9:75. https://doi.org/10.1186/1471-2164-9-75.

- Hasman H, Saputra D, Sicheritz-Ponten T, Lund O, Svendsen CA, Frimodt-Møller N, Aarestrup FM. 2014. Rapid whole-genome sequencing for detection and characterization of microorganisms directly from clinical samples. J Clin Microbiol 52:139–146. https://doi.org/10.1128/JCM.02452-13.
- Larsen MV, Cosentino S, Lukjancenko O, Saputra D, Rasmussen S, Hasman H, Sicheritz-Pontén T, Aarestrup FM, Ussery DW, Lund O. 2014. Benchmarking of methods for genomic taxonomy. J Clin Microbiol 52:1529–1539. https://doi .org/10.1128/JCM.02981-13.
- Paauw A, Caspers MPM, Schuren FHJ, Leverstein-van Hall MA, Delétoile A, Montijn RC, Verhoef J, Fluit AC. 2008. Genomic diversity within the Enterobacter cloacae complex. PLoS One 3:e3018. https://doi.org/10.1371/ journal.pone.0003018.